



Credit: Dorcas

Driving equitable action on  
NCDs and healthy ageing to  
achieve health for all at all ages

**HelpAge**  
International

## Key messages

**Non-Communicable Diseases (NCDs) and mental health and neurological conditions, including dementia, are the leading cause of death and disability globally.** Prior to COVID-19, NCDs were responsible for 74 per cent of all deaths globally and 82 per cent of years lived with disabilities.

**Population ageing is a triumph of human development, but our ability to reap its benefits is threatened by the alarming impact of NCDs** which are contributing to a growing gap between life-expectancy and healthy life expectancy, with profound implications for people, systems and societies.

**Older people are the age group most at risk from NCDs, yet are often left behind in policy and practice.** In 2021, 85 per cent of all deaths from NCDs globally were among people aged 55 and over, while rates of Years Lived with a Disability caused by NCDs were highest for older people. Despite this, and despite some acknowledgements of the disproportionate impact of NCDs on older people in the political declaration of the High Level Meeting (HLM) on NCDs in 2018, older people's needs and rights remain poorly addressed in policy and action on NCDs and health, and they face multiple barriers to accessing integrated NCD prevention and care, including barriers related to the availability, accessibility, acceptability and quality of services, and ageism and age discrimination in their funding, design and delivery. They are also the age group most at risk of catastrophic health expenditure.

**There is a strong gender dimension to NCDs, ageing and disability.** While men are likely to die earlier from NCDs, women experience a greater burden of ill health and disability from NCDs across the life-course. At the same time women of all ages – including older women – are the main providers of health and care for people living with NCDs and mental health and neurological conditions, including dementia.

**While early life interventions are critical, there is no age limit to the individual and economic benefits of prevention.** Yet older people are often excluded from action to address the determinants of NCDs.

**Opportunities are being missed for tackling NCDs and achieving better outcomes for all** due to a focus on vertical disease programming and failures to invest in the systems, services and workforce needed to deliver integrated, person-centred and community-based care that holistically responds to older people's more complex health and care needs whilst promoting healthy ageing.

**Health funding is hampering progress on NCDs and healthy ageing** by failing to keep pace with demographic and epidemiological shifts that demand greater investment in integrated NCD prevention and care, and systems that support people's intrinsic capacity and functional ability across the life-course.

**Data systems present an immediate threat to driving equitable progress on NCDs** by excluding older people through the use of ageist and discriminatory age caps, including the 'premature mortality' indicator (defined as deaths before the age of 70) in NCD frameworks, or by rendering them invisible through a lack of disaggregation.

**While the challenge of NCDs is significant, their impact is not inevitable.** By mainstreaming NCDs and mental health into progress towards universal health coverage (UHC) and primary health care (PHC), whilst investing in multi-sector 'health in all policies' approaches that tackle NCD risk factors, address health inequities and promote health and wellbeing across the life course, we can reap the benefits of healthy ageing for people, systems and societies.

## Call to action

We join with the NCD Alliance in their calls to **accelerate implementation, break down silos, mobilise investment, deliver accountability, and engage communities**. Considering older people's needs and rights within the NCD agenda, we specifically call on governments and health stakeholders at all levels to:

**Recognise and respond to the disproportionate burden of NCDs on older people and promote an equity- and rights-based approach to action at all levels.**

**Mainstream NCDs and mental health within UHC and PHC as part of efforts to reorientate health and care systems to meet the needs of older people and promote healthy ageing for all.** This must include:

Investing in age and disability inclusive and gender responsive systems that deliver equitable, integrated and person-centred care through strong primary health care approaches that engage and empower people and communities.

Including NCDs, mental health and neurological services within UHC benefit packages and ensuring people's access to the full continuum of services with financial protection – from health promotion and disease prevention to screening, early diagnosis, treatment, rehabilitation, palliative and end of life care, and long-term care and support, alongside access to the medicines, vaccines and assistive technologies needed by older people living with NCDs in all settings.

**End the discriminatory and ageist focus on 'premature mortality' within the NCD agenda** by removing upper age caps in NCD monitoring frameworks and data systems; strengthening sex, age and disability disaggregated data; and giving greater attention to NCD-related mortality, morbidity and disability across the life-course, including in research.

**Invest in the health and care workforce needed to deliver person-centred, integrated and rights-based NCD and mental health services and promote healthy ageing**, including through recognising, reducing and redistributing unpaid care, and rewarding and representing paid care, to advance the rights of women and girls of all ages, in line with the Beijing Platform for Action.

**Advance access to rights-based, integrated, person-centred and community-based services and support for people living with NCDs, mental health and neurological conditions, including dementia.** This must include accelerating de-institutionalisation and ensuring all services and support align with the Convention on the Rights of Persons with Disabilities (CRPD) and other human rights instruments.

**Integrate essential NCD and mental health services into every part of the emergency cycle** and ensure services and the workforce delivering them are able to respond to the needs of older people living with NCDs in humanitarian settings.

**Advance action on the environmental, social, economic and commercial determinants of health, including climate change, and tackle health inequities across the life-course**, adopting an intersectional, gender transformative and equity-based approach. This must include implementation of WHO "best buys".

**Engage older people living with NCDs, mental health and neurological conditions, and civil society organisations working with them, in the design and delivery of responses at all levels**, recognising them as agents of change in achieving health for all at all ages.


# Introduction

The world is ageing. By 2030, 1.4 billion people globally will be aged 60 and over, making up 17 per cent of the total population.<sup>1</sup> By 2050, this figure will rise to 2.1 billion when older people will constitute 22 per cent of the population. The majority of these older people (more than 70 per cent) live in low- and middle-income countries and this will rise to 80 per cent by 2050.

The global 'demographic transition' towards population ageing has been accompanied by an 'epidemiological transition' with all countries in the world experiencing a shift from a predominance of infectious, communicable diseases towards non-communicable diseases (NCDs). NCDs include conditions such as cancers, cardiovascular disease, stroke, chronic respiratory diseases and diabetes, as well as mental health and neurological conditions, including dementia. Prior to the COVID-19 pandemic, NCDs contributed to 74 per cent of deaths globally – an equivalent of 41 million deaths each year – and 82 per cent of all years lived with a disability (YLDs).

Population ageing is a triumph of human development, but our ability to reap the benefits of this longevity for people and societies is currently threatened by the alarming impact of NCDs which are contributing to a growing gap between life-expectancy and healthy life expectancy, especially for women.<sup>2</sup> This means that while people are living longer, they are spending a greater proportion of their lives in ill health or with a disability. (See Image 1).

**Image 1**



As with healthy ageing more broadly, NCDs are influenced by social, economic, environmental and commercial determinants of health shaping the conditions in which we are born, grow, live, work and age. These determinants interact with individual characteristics, including age, gender and disability, among others, and influence people's exposure to key NCDs risk factors - including tobacco use, physical inactivity, the harmful use of alcohol, unhealthy diets, overweight and obesity, and air pollution, all of which increase the likelihood of experiencing and dying from NCDs. These determinants also shape the extent to which people are able to benefit from health promotion initiatives and NCD prevention, early diagnosis, treatment and care. Across all settings, these inequities mean the greatest burden of NCDs, both in terms of mortality and morbidity, is faced by low- and middle-income countries and by the poorest communities - particularly those facing multiple and intersecting inequalities and living in insecure, fragile and climate affected settings.

Across age groups, older people face the greatest risk of ill health, disability and death from NCDs. Despite this, limited attention and action has been focused on the needs and rights of older people within the NCD agenda globally or addressing the unique issues they face. This gap is reinforced by the exclusion of older people from NCD monitoring frameworks and by ageism in the funding, design, delivery and reporting of health at all levels.




Addressing NCDs across the life-course – including in later life – is therefore an issue of equity. But it is also one of economics. Tackling inequalities in social determinants of health and promoting access to NCD prevention and care across the life-course as part of progress towards Universal Health Coverage (UHC) can ensure we prevent, delay and reduce ill health, disability and death caused by NCDs for people of all ages. Failing to do so, will increase pressure on health systems and undermine progress towards UHC, health equity and achievement of the Sustainable Development Goals more broadly.

As countries drive progress in commitments made in the Political Declaration on Universal Health Coverage in 2023 and look ahead to the Fourth High Level Meeting on NCDs taking place in September 2025, this briefing considers the intersection of older age with NCDs and disability, and their importance for health systems and societies alike. It outlines the unique challenges facing older people living with NCDs in low- and middle-income countries, and the barriers they face to accessing NCD prevention and care as part of integrated and person-centred health and care services that respond to their needs and rights. The briefing concludes with key actions that governments and health actors at all levels must take to drive equitable action on NCDs as part of progress towards **achieving UHC fit for an ageing world**.<sup>3</sup> This will ensure we meet the needs of older people whilst reaping the benefits of healthy ageing for individuals, systems and societies.

## Older people are the age group most at risk from NCDs

While there is great diversity in how people age, experiencing one or more NCD becomes more common in later life and older people consequently face greater risk from these conditions. In 2021, 85 per cent of all deaths from NCDs globally were among people aged 55 and over, while rates of Years Lived with a Disability caused by NCDs were highest for older people.<sup>4</sup>



Prior to COVID-19, the five leading causes of death among older people were NCDs, including cardiovascular disease, cancer, chronic respiratory diseases, neurological disorders, including dementia, and diabetes and chronic kidney disease. Analysis shows that between 1990 and 2019, while improvements in death rates among people aged 70 and over from some NCDs were seen to decrease, including from cardiovascular, chronic respiratory diseases, and some cancers, increased death rates were noted for a number of conditions, including Alzheimer's disease and other dementias (+29 per cent), lung cancer (+12 per cent), diabetes (+16 per cent), and chronic kidney disease (32 per cent).<sup>5</sup>

While globally the focus on the challenge of NCDs tends to be on mortality, critically, NCDs are also the largest cause of ill health and disability across the life course, especially in later life, with NCDs contributing to 87 per cent of all Years Lived with Disability (YLD) among people aged 55 and over in 2021. Leading causes of YLDs among older people include musculoskeletal disorders, sense organ diseases, cardiovascular diseases, diabetes and chronic kidney diseases, neurological disorders, including dementia, and mental health conditions.<sup>6</sup>

# Mental health and neurological conditions have a profound and growing impact on older people

Mental health conditions have a profound impact on older people but often receive limited attention. According to the World Health Organization, around 14 per cent of adults aged 60 and over live with a mental health condition.<sup>7</sup> These conditions account for 10 per cent of all Years Lived with Disability among older people, while over a quarter (27.2 per cent) of deaths from suicide globally are among people aged 60 or over.<sup>8,9</sup> The most common mental health conditions for older adults are depression and anxiety which older people may be at increased risk of because of higher incidence of poverty, social isolation and loneliness, bereavement and other life changes, physical or neurological health conditions, or lack of access to quality support and services.<sup>10</sup> This is particularly the case for older people living in humanitarian settings (see Box 2 below).

Globally, neurological conditions, including conditions such as stroke, Parkinson's and dementia, are now the leading cause of ill health and disability worldwide.<sup>11</sup> The prevalence of most neurological diseases increases sharply with age, and they are often associated with more intense health and social care needs. This is particularly the case for Alzheimer's Disease and other dementias. Around 57 million people have dementia worldwide and in 2019, dementia was estimated to cost economies globally 1.3 trillion US dollars, with approximately 50 per cent of these costs attributable to care and support provided by informal caregivers – mostly women of all ages – delivering on average, five hours of care and support per day.<sup>12</sup>

Older people living with mental health or neurological conditions such as dementia, and those living with psychosocial disabilities, are at high risk of experiencing violations of their human rights. An over-reliance on biomedical approaches to treatment options and inpatient care, and a failure to promote and invest in community-based, integrated, person-centred, gender-responsive and rights-based models of health, rehabilitation, palliative, end of life and long-term care and support, leads to these groups experiencing stigma, discrimination and routinely being denied their right to dignity, autonomy, independence, choice and participation, and freedom from violence, abuse and neglect<sup>13,14,15</sup>

## The majority of older people are living with more than one condition

As NCDs increase with age, older people are also at greater risk of experiencing two or more conditions at the same time, with more than half (51 per cent) of older people living in the community globally estimated to be living with two or more long-term conditions ('co-' or 'multi-' 'morbidity').<sup>16</sup> Multimorbidity is associated with greater and more complex health and care needs and poorer levels of functional ability, quality of life and higher rates of mortality.

NCDs also often increase the risk someone faces from infectious diseases. For example, people living with NCDs are at heightened risk of respiratory illnesses. During COVID-19, an estimated 60 to 90 per cent of mortality in COVID-19 cases was attributable to the presence of either one or more NCD.<sup>17</sup> Certain NCDs are also associated with even higher risk from respiratory illnesses. For example, people with diabetes face a two-to-four-fold higher risk of active Tuberculosis (TB) and up to 30 per cent of individuals with TB are estimated to have diabetes.<sup>18</sup>

The interaction between NCDs and HIV/AIDS is also a critical area for health and care but often overlooked. With increasing access to antiretroviral therapy for HIV and the growth of new infections amongst older people, an “ageing” of the epidemic is now occurring. UNAIDS estimates that the number of people aged 50 years or older with HIV infection globally increased from 5.4 million in 2015 to 8.1 million in 2020.<sup>19,20,21</sup> Older people with HIV have higher levels of multimorbidity compared with people of similar age without HIV,<sup>22</sup> including a higher frequency of NCDs and the need for multiple treatments and medications (‘polypharmacy’).<sup>23</sup> While a considerable research gap exists in this area, emerging evidence illustrates how the two epidemics of NCDs and HIV interact with one another within a context of poverty, inequality and inequitable access to healthcare resulting in compounded challenges for older people living with HIV and NCDs, including experience of stigma and discrimination.<sup>24</sup>

## There are important gender differences to NCDs in later life

Globally, men outnumber women until age 50, after which women predominate due to longer life expectancies. By 2030, 54 per cent of the world’s 1.4 billion older people will be women, rising to 60 per cent among people aged 80 and over.<sup>25</sup> However, partly due to their longer life expectancies, women spend more of their lives in ill health or with disability, including more years with NCD-related disability and a higher likelihood of co- or multi-morbidity.<sup>26,27</sup>

Gender plays a key role in shaping social determinants of health and exposure to NCD risk factors and access to services. Inequalities experienced women and girls across the life-course lead to them facing higher risk of multi-dimensional poverty than men, with significant impact on their health and wellbeing.<sup>28</sup> Women and girls also often receive less education than men, limiting their capacity to inform and protect themselves against risk factors for NCDs, including unhealthy diets, **air pollution, tobacco and alcohol use**. In some settings, women present lower levels of physical activity compared to men as a result of social and cultural customs related to gender and mobility, and are also at greater risk of indoor air pollution due to household tasks.<sup>29,30</sup> In some settings, women present lower levels of physical activity compared to men as a result of social and cultural customs related to gender and mobility, and are also at greater risk of indoor air pollution due to household tasks.<sup>31,32</sup>

Gender differences result in older women being at greater risk of certain NCDs, including heart disease, risk of stroke and osteoporosis, cervical and breast cancer, and risks and complications related to their sexual and reproductive health, including menopause and post-menopause.<sup>33</sup> Women are also disproportionately affected by Alzheimer’s disease and other dementias, experiencing higher morbidity and mortality due to dementia than men.<sup>34</sup>

At the same time, women of all ages – including older women – are the main providers of health and social care for those living with NCDs and other health conditions, both formally and informally.<sup>35</sup> As many as 70 per cent of the global healthcare workforce is female, while women and girls are estimated to do at least two and a half times more unpaid household and care work than men, including care for older people.<sup>36</sup> This care need/care provision dynamic can place older women in a precarious position, compounded by the effects of power imbalances and gender inequalities experienced by women and girls across their life-course and by discrimination they face on the grounds of gender, age, disability, and other characteristics in older age.<sup>37</sup> These factors put them at higher risk of exclusion and of experiencing violence, abuse and neglect,<sup>38</sup> with profound impacts on their physical and mental health and wellbeing, and on their access to services and support that meets their needs.

# There is no age limit to the benefits of prevention

Despite their significant impact, many of the consequences of NCDs are avoidable. Acting on the determinants and risk factors for NCDs across the life-course, and inequalities in people's exposure to these, whilst ensuring access to community based - integrated NCD prevention and care as part of progress towards UHC, offers the opportunity to prevent, reduce and delay more acute health and care needs and many deaths.

While the role of health promotion and disease prevention from an early age is critical, the benefits of prevention extend across the life-course, with continued potential for significant positive impact in later life. In 2019, 58 per cent of Disability Adjusted Life Years – that is, the sum of the years of life lost due to premature mortality and the years lived with a disability – were attributable to risk factors in people aged 70 and over.<sup>39</sup> The top five risk factors were high blood pressure, high blood sugar, smoking, high cholesterol, and overweight/obesity, many of which are influenced by modifiable behaviours, such as tobacco use, physical inactivity, unhealthy diet, exposure to air pollution and the harmful use of alcohol.<sup>40</sup>

An example is the case of diabetes. In 2021, an estimated 537 million people aged 20–79 globally had diabetes with prevalence increasing with age. This number is projected to reach 783 million by 2045.<sup>41</sup> Type 2 diabetes, which accounts for the majority of the cases, can lead to multiple organ complications and heightened risk of a range of other health issues and complications. Yet by addressing modifiable risk factors, it is possible to prevent and/or delay the onset of Type 2 diabetes and its complications. Despite this, an estimated 240 million people globally are living with undiagnosed diabetes, meaning almost one-in-two adults with diabetes are unaware they have the condition. Almost 90 per cent of people with undiagnosed diabetes live in low- and middle-income countries, with more than half of people living with diabetes in Africa, South-East Asia and the Western Pacific undiagnosed.

To drive progress, countries must accelerate equitable action on the major NCDs and their leading risk factors, in line with WHO's five-by-five approach and evidence-based and cost-effective interventions outlined in WHO's "Best Buys".<sup>42</sup> This must include investing in primary health care approaches that prioritise prevention and promote healthy ageing across the life-course within progress towards UHC, recognising this as an investment rather than a cost. To reap these benefits, communities must be engaged and empowered in their own health and wellbeing (see Box 1) and barriers older people face to accessing integrated and person-centred NCD prevention, screening, early diagnosis and care must be addressed.



# Older people face multiple barriers to accessing integrated and person-centred NCD prevention and care


Despite older people being the age group most at risk from NCDs, they are often furthest behind in accessing NCD prevention and care. For example, while people living with NCDs are, in general, more likely to experience catastrophic and impoverishing health spending than those without NCDs, older people face the greatest risk.<sup>43</sup> (See Image 2).<sup>44</sup>

In many settings, limited funding for NCDs, mental health, and for services that holistically meet the needs of older people, means that opportunities are being missed for effectively preventing

and managing NCDs and promote healthy ageing. Countries have demonstrated almost no progress since 2000 in expanding service capacity and access to prevention, screening, early diagnosis and treatment for NCDs.<sup>45</sup> Only 54 per cent of 194 countries surveyed by WHO in 2021 reported general availability of 11 essential NCD medicines, with one in five countries reporting that only six or fewer were generally available.<sup>46</sup>

Access to rehabilitation, assistive technologies (AT), palliative care, end of life care and long-term care and support are also critical gaps related to meeting the needs and upholding the rights of older people living with NCDs, mental health and neurological conditions, and promoting healthy ageing. WHO estimates that 2.5 billion people – or 1 in 3 people – need one or more assistive products, rising to two thirds (69 per cent) of older people, but that nearly one billion people of all ages are denied access.<sup>47</sup> This is particularly the case in low- and middle-income countries, where, on average, only 10 per cent for those who need AT are able to access it.<sup>48</sup> Over 56.8 million people are estimated to require palliative care, with NCDs account for almost 69 per cent of adult need.<sup>49</sup> Yet worldwide, only about 14 per cent of people who need palliative care currently receive it.<sup>50</sup>

**Image 2: Percentage of population experiencing catastrophic health expenditure**



**Only 54%**

of 194 countries surveyed by WHO report general availability of 11 essential medicines for NCDs which are more common in later life.



Access to assistive technology for those who need it varies globally and is as


**low as 3%**

in some countries.

The failure to promote disability inclusion within health systems, and in action on NCDs specifically, in line with WHO's Global report on health equity for persons with disabilities, is another major barrier to older people living with NCDs and mental health and neurological conditions accessing the services and support they need.<sup>51</sup> Considering the strong interplay between ageing, NCDs and disability, with both older people and persons with disabilities facing greater risk of NCDs, and NCDs being the leading cause of years lived with disability

globally, this failure exacerbates health inequities and contributes to worse outcomes for those living with NCDs of all ages.

Even when services are available and accessible, older people often report that ageism and age discrimination affect their right to access them on an equal basis with others. This includes assumptions from health care workers that NCDs are just a normal part of ageing.<sup>52</sup>



More generally, a focus on vertical disease programming and a failure to invest in the systems, services and workforce needed to deliver integrated, person-centred and community-based models of care that holistically respond to older people's more complex health and care needs whilst promoting healthy ageing for all, means opportunities for addressing NCDs and promoting health and wellbeing across the life-course are being missed. As countries progress towards UHC, it is critical to address this gap and mainstream NCD prevention and care, mental health and neurological conditions, and interventions that promote healthy ageing more broadly, within UHC benefit packages and all health system building blocks.

### **Box 1: Scaling up NCD prevention and care through primary health care interventions engaging older people and their communities**

Scaling up Non-Communicable Disease Interventions in Southeast Asia (SUNI-SEA) was an action research project implemented through a consortium of partners including HelpAge in Vietnam and Myanmar between 2019-2023. It highlighted the value of engaging people and communities to drive progress on NCDs.

Community NCD interventions were implemented by community-based organisations with the support of volunteers in Vietnam and Myanmar. Following skills building, the volunteers conducted community level screening for NCDs, planned and facilitated health promotion sessions, motivated people for peer support and self-care activities, referred people at risk of NCDs to health facilities and monitored, reported and evaluated their activities.

In Vietnam, the project activities were implemented at primary health care facility level by government partners, in-country research institutes and older-people led Intergenerational Self-Help Groups. Activities included strengthening the quality of NCD care and treatment services through health centre management training, capacity building, updated service guidelines, development of user-friendly job aids, supportive supervision, and strengthening the linkages between the community and local health facilities. Digital technology was used to support stronger community-based NCD screening, management and data and 295 health volunteers were trained in best practice screening and health promotion alongside 126

commune-health staff who strengthened their capacities around screening, diagnosis and treatment for NCDs.

In Myanmar, where the context was different, 30 health volunteers were trained and a self-screening mobile application tool was used to reach close to 2000 people who are now able to regularly screen for diabetes, hypertension and cardiovascular disease, alongside common mental health conditions.

Through its activity, the consortia demonstrated that community interventions, led by older people, boost self-care and increase awareness about NCDs, underlining that adopting a community-based approach, integrated with strong primary health care, is key to achieving UHC and improving health outcomes. To work well, community health services must be community-led, people-centred, and delivered through co-ownership. They must be gender responsive and ensure the meaningful engagement of diverse communities. Finally, sustainable scaling up requires engaging local authorities and advocating for bottom-up funding to support implementation. Read more [here](#).

## Health funding is hampering progress on NCDs and healthy ageing

The barriers older people face to accessing services that meet their needs are compounded by ageism and age discrimination in health financing. This is seen in the limited funding for older people's health and wellbeing, as well as in NCD financing.

At global level, analysis of Development Assistance for Health (DAH) from 2017, for example, found that DAH targets younger more than older age groups relative to their disease burden, with 90 percent of DAH going to people younger than 60.<sup>53</sup> Rather than responding to population ageing, this trend was seen to be increasing, with analysis finding that diseases causing health burden at older ages were actually deprioritised between 1990 and 2013, despite the demographic and epidemiological changes that increased their importance.<sup>54</sup> Meanwhile, funding for NCDs received just 2.3 per cent of funding in 2023, despite being the leading cause of disease, disability and death globally.<sup>55</sup>

More broadly, the focus of DAH on single, largely infectious disease funding is also missing opportunities for investing in health system strengthening and integrated primary health care approaches that would benefit people living with NCDs across the life course.<sup>56</sup>

At national level, despite the evidence of the devastating impact of NCDs and mental health, and the existence of a strong evidence base of proven cost-effective solutions, governments have been slow to invest in NCD prevention and care, mental health and wellbeing, and healthy ageing more broadly.<sup>57</sup> This failure continues to lead to dire outcomes and missed opportunities for individuals, health and care systems, and societies. Governments urgently need to mobilise additional domestic revenues for action, including through investing in WHO's best buys for NCDs which are expected to yield a return of 7:1, while other studies estimate returns of 19:1 for investments in NCDs.<sup>58,59</sup> NCDs and mental health must be integrated into UHC and PHC, and health financing policies and mechanisms must promote prevention, advance access to integrated and person-centred care that supports healthy ageing, and protect people from catastrophic and impoverishing health spending.<sup>60</sup> To support greater public resources for health in lower income countries, there is also urgent need to create a fairer global economic system that would allow the poorest countries to invest sufficiently in health.<sup>61</sup>

# Data systems present an immediate threat to driving equitable progress on NCDs

Despite facing the greatest risk from NCDs, older people are excluded from official statistics on NCDs at local, national and global levels or rendered invisible through a lack of age, sex and disability disaggregated data.

WHO's NCD Global Monitoring Framework uses an age-bracketed indicator on unconditional probability of dying from four main NCDs, limited to the ages of 30-70. This indicator is also used to inform progress towards SDG indicator 3.4 which focuses on 'reducing by one third premature mortality from non-communicable diseases'. The NCD Global Monitoring Framework also uses age brackets in other indicators, including the indicator on cervical cancer screening which is limited to the ages 30-49, despite cervical cancer continuing to occur in older women. The framework's focus on 'premature mortality' is discriminatory and ageist, whilst also failing to capture the impacts of NCDs beyond mortality, including both morbidity and disability related to NCDs.

Another critical gap is the WHO STEPwise approach to surveillance (WHO STEPS), a survey mechanism for collecting national level data on risk factors for NCDs which is also used to monitor global level progress. Again, despite the prevalence of NCDs amongst people in older age, WHO STEPs typically only includes people between the ages of 18-69, while in a number of countries the cut off is even lower. For example, analysis by HelpAge in 2017 found that only six of around 40 countries in Africa that had conducted a STEPS survey had included people over the age of 64 in their most recent survey.<sup>62</sup> Analysis of the Global Burden of Disease study data, meanwhile, highlighted that coverage of risk factor data for the population aged 70 and over actually decreased in almost 30 per cent of the GBD locations between 1990 and 2019, while since 1990 no information is available for nine risk factors in older adults.<sup>63</sup>

Older people are also excluded from wider data systems that inform NCD policy and practice. For example, current measures of UHC, including the 'access' indicator (3.8.1) in the SDG indicator framework, do not include indicators such as physical access to health facilities, or staff skills, knowledge and attitudes – factors that are critical to understanding the barriers faced by older people. There are also shortcomings in the data sources typically used to populate these indicators, including WHO STEPs mentioned above and the Demographic and Health Survey (DHS), which typically excludes women over the age of 49 and men over the age of 59, rendering older people with and without NCDs invisible.

Even where data is collected on older age groups, systems often fail to analyse, report and use sufficiently disaggregated data for understanding the diversity of older people's health and care needs and preferences, including disaggregation by sex, gender, age, disability, location and socio-economic group to measure inequalities and inequities in relation to health status, access to services and outcomes.

Such data challenges pose one of the greatest threats to measuring and driving equitable progress on NCDs. The indicators also serve to reinforce and exacerbate age discrimination in NCD policy, practice, research and funding from global to local levels and must be addressed as an urgent matter of human rights.

## **Box 2: Older people living with NCDs in humanitarian settings**

A large and rapidly growing number of older people are affected by humanitarian crises, in which they are often among the most at risk, yet most overlooked. Globally the proportion of the population aged 50 and over in fragile countries, where humanitarian emergencies are more likely to occur, is expected to rise from 12.3 per cent (219.9 million) in 2020 to 19.2 per cent (586.3 million) in 2050.<sup>64</sup>

Humanitarian emergencies include natural disasters, armed conflict, outbreaks and other health emergencies, and may combine multiple crises, such as war and the ever-increasing threat from climate change which presents significant and growing risk to older people living with NCDs who are more at risk from its consequences.<sup>65</sup> Over time, an acute emergency can become protracted, with people potentially displaced from their homes for decades.

Despite their high levels of need, older people in humanitarian emergencies often find themselves systematically excluded. In addition to the barriers older people face in general to accessing the services they need, natural disasters, conflict, climate or other crises can cause disruptions in access to existing services and support and impair the capacity of systems to meet people's essential needs.

In some settings, health systems and services that were previously provided within a country may be seriously undermined or completely destroyed, along with access to other goods and services that people living with NCDs may rely on to manage their condition or conditions, including medicines, foods, assistive technologies or other products, increasing the risk of life-threatening complications. The impact of emergencies and disruption in care also has a profound impact on people's mental health and psychosocial wellbeing further exacerbating existing conditions.

While attention to NCDs in humanitarian response is increasing, including through the development of the WHO NCD kit for emergency settings and processes such as the high-level technical meeting on NCDs in humanitarian settings in 2024, there is urgent need to accelerate progress.<sup>66</sup>

Older people's needs and rights must be addressed within these efforts, as one of the groups at most risk. This means ensuring age and disability inclusive and gender responsive approaches are promoted throughout humanitarian preparedness and response, in line with the Humanitarian Inclusion Standards, including by ensuring that humanitarian actors at all levels are trained in how to respond to the needs and rights of older people in general and those living with NCDs and multimorbidity specifically.

To support this, NCD prevention and care in humanitarian settings must move away from disease specific interventions towards more integrated approaches that better respond to older people's more complex health and care needs. This includes their need not only for NCD prevention and disease focused management but also for rehabilitation, palliative care, and long-term care and support which must be recognised as essential components of NCD services in humanitarian settings. It is also critical to ensure mental and psychosocial health are fully integrated into emergency preparedness and response at all levels.

Older people living with NCDs, caregivers and communities must be engaged and empowered at all stages of emergency preparedness and response, to ensure their experiences shape the design and delivery of activities. This must include strengthening the collection, analysis, reporting and use of sex, gender, age and disability disaggregated data on people of all ages to inform equity-based responses.



# A call to action: tackling NCDs and promoting healthy ageing for all

Three political declarations on NCDs have committed Heads of States and governments to provide strategic leadership and to scale up action for the prevention and control of NCDs and the promotion of mental health, recognising them as major challenges for the health and well-being of all people and more broadly for sustainable development. At the High-Level Meeting on UHC in 2023, Heads of States and Governments also committed to promote and implement policy, legislative, regulatory and fiscal measures to minimise the exposure to main risk factors of NCDs and to scale up efforts in primary and specialised health services for NCD prevention, screening, treatment and control.

Yet today, despite significant policy progress and strong leadership in some countries, implementation has lagged behind.<sup>67</sup> In 2024, the world is off track to meet the global NCD targets that are set to expire in 2025 and 2030, with many promises made at the last High-Level Meeting in 2018 unmet. COVID-19 has put the response even further off-track with the greatest impact on those most at risk, including older people living with NCDs and mental health and neurological conditions.

The NCD Alliance highlights that projected figures for NCD prevalence are even more cause for concern.<sup>68</sup> Health systems already struggling to handle the NCD burden are unprepared for the future. The number of people living with diabetes is expected to more than double globally by 2050, to at least 1.3 billion. Cancer too will double, with 35 million new cases per year foreseen by 2050. Cardiovascular diseases accounts for 18 million deaths per year – this figure will reach 23 million by 2030. Chronic kidney disease is increasing worldwide at a rate of 8 per cent per year; by 2040, it is projected to be the fifth highest cause of death. And the number of people living with dementia is projected to nearly double every 20 years, reaching 139 million people by 2050.

Despite these figures and the ever-increasing challenge of NCDs, multisectoral action plans to address their causes and consequences are still lacking in roughly half of countries, and despite some improvement, only 57 per cent of countries have set time-bound national NCD targets and indicators. Equally, despite commitments made within the UN political declarations on UHC in 2019 and 2023, in many places, NCDs are failing to be integrated into action on UHC, leaving behind an estimated 20 per cent of the global population living with NCDs.<sup>69</sup> This is especially true for those who are at highest risk, including older people.

The disproportionate burden of NCDs on older people was recognised in the political declaration of the third HLM on NCDs in 2018, while the unique challenges older people face in relation to accessing health and care services that meet their needs were highlighted in the political declarations of the HLMs on UHC in 2019 and 2023. However, there remains a failure to translate these acknowledgments into focused attention to older people within the global NCD or health agendas and in funding, policy and action at national levels. This gap is reinforced by older people's continued exclusion from data systems.

These failures clearly reflect yet another example of ageism and age discrimination in development, despite a focus on leaving no one behind and reaching the furthest behind first. But they represent not just a failure of equity, but also of economics. Analysis by the World Bank found that 77 percent and 75 percent deaths of those aged 40–59 and 60–79 respectively were avoidable in 2019, alongside 88 percent of deaths in those aged 20–39.<sup>70</sup> Estimates for Latin American countries shows one of the highest levels of avoidable mortality in older age groups, with heart disease leading the ranking of the prevalence of chronic diseases that are responsible for most of these deaths, alongside respiratory infections. Their

analysis estimates that investing in a minimal “starter” package of NCD interventions, could contribute to at least 150 million deaths across all LMICs being avoided by 2050, including those among older people, and about 8 million in 2050 alone. They estimate the economic value of this avoidable mortality at over US\$3.2 trillion in 2050.

As countries drive progress in commitments made in SDGs, the Political Declarations on Universal Health Coverage in 2023 and Political Declarations on NCDs in 2018, and look ahead to the Fourth High Level Meeting on NCDs and mental health and wellbeing, we need transformative leadership to drive progress for people of all ages and reap the associated benefits for individuals, systems and societies. Action must place those most at risk - including older people - at the centre and must simultaneously address the social, economic, environmental and commercial determinants of NCDs, including climate change, and invest in prevention, while meeting the needs and upholding the rights of those living with NCDs, mental health and neurological conditions.

We join with the NCD Alliance in their calls to **accelerate implementation, break down silos, mobilise investment, deliver accountability, and engage communities.**

Considering older people’s needs and rights within the NCD agenda, we also call on governments and health stakeholders at all levels to:

- Recognise and respond to the disproportionate burden of NCDs on older people and promote an equity- and rights-based approach to action at all levels. This must ensure older people’s needs and rights are at the centre of the NCD and mental health agendas and that equity guides the funding, design and delivery of action.
- Mainstream NCDs and mental health within UHC and PHC as part of efforts to ensure health systems meet the needs of older people and promote healthy ageing for all. This must include:
- Investing in age- and disability-inclusive and gender-responsive health and care systems that deliver equitable, integrated and person-centred care through strong primary health care approaches that engage and empower people and communities.
- Including NCDs, mental health and neurological services within UHC benefit packages and ensuring people’s access to the full continuum of services with financial protection – from health promotion and disease prevention to screening, early diagnosis, treatment, rehabilitation, palliative care, end of life care and long-term care and support, alongside access to the medicines, vaccines and assistive technologies needed by older people living with NCDs in all settings.
- End the discriminatory and ageist focus on ‘premature mortality’ within the NCD agenda by removing upper age caps in NCD monitoring frameworks and data systems; strengthening the collection, analysis, reporting and use of age, sex and disability disaggregated data on people of all ages; and giving greater focus to measuring and addressing morbidity and disability caused by NCDs among people of all ages. This should be supported by greater resources for research on NCD mortality, morbidity and disability in later life and effective interventions for addressing these.
- Invest in the health and care workforce needed to deliver person-centred, integrated and rights-based NCD and mental health services and promote healthy ageing, and supported by community-based approaches ensuring a well-paid, well-trained, well-resourced, multidisciplinary and gender equal workforce able to respond effectively and holistically to the diverse and more complex health and care needs of ageing populations. This must include recognising, reducing, and redistributing unpaid care and rewarding and representing paid care to advance the rights of women and girls of all ages, in line with the Beijing Platform for Action.

- Advance access to rights-based, person-centred and community-based services and support for those living with mental health and neurological conditions, including dementia. This must include accelerating de-institutionalisation and ensuring all services and support align with the Convention on the Rights of Persons with Disabilities (CRPD) and other human rights instruments.
- Integrate essential NCD and mental health services into every part of the emergency cycle and ensure services and the workforce delivering them are able to respond to the unique needs of older people living with NCDs, including the need for inclusive, person-centred and integrated health and care services across preparedness and disaster risk reduction, through the immediate emergency response.
- Commit to advance action on the environmental, social, economic and commercial determinants of health, including climate change, and tackle health inequities across the life-course, adopting an intersectional, gender-transformative and equity-based approach. This must include implementation of WHO's "Best Buys".
- Engage older people living with NCDs, mental health and neurological conditions, and civil society organisations working with them, in the design and delivery of responses at all levels, recognising them as agents of change in achieving health for all at all ages.

## Endnotes

<sup>1</sup> United Nations Department of Economic and Social Affairs, Population Division (2022), *World Population Prospects 2022*. POP/03-1: Total population (both sexes combined) by select age group, region, subregion and country, annually for 1950-2100 (thousands), Medium variant, Online edition.

<sup>2</sup> World Health Organization (WHO), *Global Health Estimates: Life expectancy and healthy life expectancy 2021*

<sup>3</sup> HelpAge International, 2024, *Achieving Universal Health Coverage fit for an ageing world*.

<sup>4</sup> Institute for Health Metrics and Evaluation (IHME), *Global Burden of Disease (GBD) 2021, Results Tool*, [vizhub.healthdata.org/gbd-results/result/989fb4795f2407d0820617c7678126e4](https://vizhub.healthdata.org/gbd-results/result/989fb4795f2407d0820617c7678126e4) (15 May 2025)

<sup>5</sup> Global, regional, and national burden of diseases and injuries for adults 70 years and older: systematic analysis for the Global Burden of Disease 2019 Study *BMJ* 2022; 376 :e068208 doi:10.1136/bmj-2021-068208

<sup>6</sup> Ibid.

<sup>7</sup> WHO, *Mental health of older adults fact sheet*. Accessed on 13.10.2024.

<sup>8</sup> IHME, *Global Burden of Disease 2021, GBD results VizHub - GBD Results* (healthdata.org) Accessed on 13.10.2024

<sup>9</sup> WHO 2019, 'Global health estimates: Summary tables: Death by cause, age and sex'. Geneva, Switzerland <https://www.who.int/data/gho/data/themes/mortality-and-global-health-estimates>

<sup>10</sup> WHO 2021, *Social isolation and loneliness among older people: advocacy brief*. Geneva: World Health Organization; 2021. Licence: CC BY-NC-SA 3.0 IGO.

<sup>11</sup> WHO, 'Over 1 in 3 people affected by neurological conditions, the leading cause of illness and disability worldwide'.

<sup>12</sup> WHO, *Dementia fact sheet*. Accessed on 10.09.2024.

<sup>13</sup> The Lancet Healthy Longevity. Ageing with HIV. *Lancet Healthy Longev.* 2022 Mar;3(3):e119. doi: 10.1016/S2666-7568(22)00041-1. Epub 2022 Feb 23. PMID: 36098283.

<sup>14</sup> Belaunzaran-Zamudio, P. F., Caro-Vega, Y., Giganti, M. J., Castilho, J. L., Crabtree-Ramirez, B. E., Shepherd, B. E., . . . Sierra-Madero, J. G. (2020). Frequency of noncommunicable diseases in people

---

50 years of age and older receiving HIV care in Latin America. PLoS One, 15(6), e0233965. <https://doi.org/10.1371/journal.pone.0233965>

<sup>15</sup> Goldstein D, Kiplagat J, Taderera C, Whitehouse ER, Chimbetete C, Kimaiyo S, Urasa S, Paddick SM, Godfrey C. Person-centred care for older adults living with HIV in sub-Saharan Africa. *Lancet HIV*. 2024 Aug;11(8):e552-e560. doi: 10.1016/S2352-3018(24)00123-1. Epub 2024 Jul 9. PMID: 38996592.

<sup>16</sup> Saifur Rahman Chowdhury, Dipak Chandra Das, Tachlima Chowdhury Sunna, Joseph Beyene, Ahmed Hossain, 'Global and regional prevalence of multimorbidity in the adult population in community settings: a systematic review and meta-analysis', *Lancet eClinical Medicine*, VOLUME 57, 101860, March 2023

<sup>17</sup> Palmer K, Monaco A, Kivipelto M, Onder G, Maggi S, Michel JP, Prieto R, Sykara G, Donde S. The potential long-term impact of the COVID-19 outbreak on patients with non-communicable diseases in Europe: consequences for healthy ageing. *Aging Clin Exp Res*. 2020 Jul;32(7):1189-1194. doi: 10.1007/s40520-020-01601-4. Epub 2020 May 26. PMID: 32458356; PMCID: PMC7248450.

<sup>18</sup> Krishna S, Jacob JJ. Diabetes Mellitus and Tuberculosis. [Updated 2021 Apr 18]. In: Feingold KR, Anawalt B, Blackman MR, et al., editors. *Endotext* [Internet]. South Dartmouth (MA): MDTText.com, Inc.; 2000-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK570126/>

<sup>19</sup> The Lancet Healthy Longevity. Ageing with HIV. *Lancet Healthy Longev*. 2022 Mar;3(3):e119. doi: 10.1016/S2666-7568(22)00041-1. Epub 2022 Feb 23. PMID: 36098283.

<sup>20</sup> Belaunzaran-Zamudio, P. F., Caro-Vega, Y., Giganti, M. J., Castilho, J. L., Crabtree-Ramirez, B. E., Shepherd, B. E., . . . Sierra-Madero, J. G. (2020). Frequency of noncommunicable diseases in people 50 years of age and older receiving HIV care in Latin America. PLoS One, 15(6), e0233965. <https://doi.org/10.1371/journal.pone.0233965>

<sup>21</sup> Goldstein D, Kiplagat J, Taderera C, Whitehouse ER, Chimbetete C, Kimaiyo S, Urasa S, Paddick SM, Godfrey C. Person-centred care for older adults living with HIV in sub-Saharan Africa. *Lancet HIV*. 2024 Aug;11(8):e552-e560. doi: 10.1016/S2352-3018(24)00123-1. Epub 2024 Jul 9. PMID: 38996592.

<sup>22</sup> Ibid.

<sup>23</sup> Belaunzaran-Zamudio et al. 2020.

<sup>24</sup> Knight L, Schatz E, Mukumbang FC. "I attend at Vanguard and I attend here as well": barriers to accessing healthcare services among older South Africans with HIV and non-communicable diseases. *Int J Equity Health*. 2018 Sep 18;17(1):147. doi: 10.1186/s12939-018-0863-4. PMID: 30227859; PMCID: PMC6145370.

<sup>25</sup> UNDESA, 2022, *World Population Prospects 2022*. POP/03-1: Total population (both sexes combined) by select age group, region, subregion and country, annually for 1950-2100 (thousands), Medium variant, Online edition.

<sup>26</sup> IHME, GBD2019.

<sup>27</sup> Saifur Rahman Chowdhury et al, 2023

<sup>28</sup> UN Women, Older women: Inequality at the intersection of age and gender, UN Women data hub, 2022. Quoted in HelpAge 2024.

<sup>29</sup> The NCD Alliance and The George Institute for Global Health, 2019, Delivering healthy lives and well-being for women and girls: Non-communicable diseases and Universal Health Coverage.

<sup>30</sup> Verma A, Gudi N, Yadav UN, Roy MP, Mahmood A, Nagaraja R, Nayak P. Prevalence of COPD among population above 30 years in India: A systematic review and meta-analysis. *J Glob Health*. 2021. Aug 21;11:04038. doi: 10.7189/jogh.11.04038. PMID: 34484706; PMCID: PMC8397327

<sup>31</sup> The NCD Alliance and The George Institute for Global Health, 2019, Delivering healthy lives and well-being for women and girls: Non-communicable diseases and Universal Health Coverage.

<sup>32</sup> Verma A, Gudi N, Yadav UN, Roy MP, Mahmood A, Nagaraja R, Nayak P. Prevalence of COPD among population above 30 years in India: A systematic review and meta-analysis. *J Glob Health*. 2021. Aug 21;11:04038. doi: 10.7189/jogh.11.04038. PMID: 34484706; PMCID: PMC8397327

- 
- <sup>33</sup> HelpAge International 2023, Achieving gender transformative UHC fit for an ageing world. Available at <https://www.helpage.org/wp-content/uploads/2023/07/Achieving-gender-transformative-UHC.pdf>
- <sup>34</sup> WHO, Dementia fact sheet. Accessed 17.07.2024.
- <sup>35</sup> WHO, Value gender and equity in the global health workforce, accessed November 2022. Quoted in HelpAge 2024.
- <sup>36</sup> Oxfam International, Not all gaps are created equal: the true value of care work, accessed November 2022. Quoted in HelpAge 2024.
- <sup>37</sup> UN Women, Older women: Inequality at the intersection of age and gender, UN Women data hub, 2022. Quoted in HelpAge 2024.
- <sup>38</sup> WHO, 2024, 'WHO calls for greater attention to violence against women with disabilities and older women'. News article, 27.03.2024.
- <sup>39</sup> Saifur Rahman Chowdhury et al, 2023.
- <sup>40</sup> Ibid.
- <sup>41</sup> WHO, Diabetes: Fact Sheet, Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes>. Accessed on 26.07.2024
- <sup>42</sup> WHO, Noncommunicable Diseases, Rehabilitation and Disability factsheet. Accessed 20.10.2024.
- <sup>43</sup> NCD Alliance and The George Institute, 2023, Paying the Price: A deep dive into the household economic burden of care experienced by people living with noncommunicable diseases.
- <sup>44</sup> WHO and World Bank, 2023, Tracking universal health coverage 2023 Global Monitoring Report.
- <sup>45</sup> WHO, Noncommunicable Diseases, Rehabilitation and Disability factsheet. Accessed 20.10.2024.
- <sup>46</sup> WHO, 2023, Assessing national capacity for the prevention and control of noncommunicable diseases: report of the 2021 global survey. Geneva: World Health Organization; 2023. Licence: CC BY-NC-SA 3.0 IGO
- <sup>47</sup> WHO and United Nations Children's Fund (UNICEF), 2022, Global report on assistive technology.
- <sup>48</sup> Ibid.
- <sup>49</sup> World Hospice and Palliative Care Alliance (WHPCA) Global Atlas of Palliative Care, 2nd Ed 2020.
- <sup>50</sup> WHO, Palliative care fact sheet. Accessed 17.10.2024
- <sup>51</sup> WHO, 2022, Global report on health equity for persons with disabilities.
- <sup>52</sup> HelpAge 2023, Healthy ageing for us all: what older people say about their right to health and health services.
- <sup>53</sup> Skirbekk V, Ottersen T, Hamavid H, Sadat N, Dieleman JL. 'Vast majority of development assistance for health funds target those below age sixty', *Health Affairs*, 2017 May; 36(5):926-930. doi: 10.1377/hlthaff.2016.1370
- <sup>54</sup> Ibid.
- <sup>55</sup> Institute for Health Metrics and Evaluation (IHME). Financing Global Health. Seattle, WA: IHME, University of Washington, 2023. Available from <http://vizhub.healthdata.org/fgh/>(link is external). Accessed March 11 2024.
- <sup>56</sup> Jonathan E. Cohen (2023) 'How Can US Global Health Assistance Adapt to Population Aging?' Online article, Just Security. November 2023. How Can US Global Health Assistance Adapt to Population Aging? ([justsecurity.org](https://www.justsecurity.org))
- <sup>57</sup> World Health Organization (2024) 'International dialogue on sustainable financing for noncommunicable diseases and mental health: Health expenditures on noncommunicable diseases and mental health: What can health accounts tell us? Technical background paper #1: Our analysis of the Health Accounts landscape and data availability has revealed substantial weaknesses". Available at



---

[https://cdn.who.int/media/docs/default-source/ncds/paper-1-nha-summary-brief.pdf?sfvrsn=2ae4bcce\\_3&utm](https://cdn.who.int/media/docs/default-source/ncds/paper-1-nha-summary-brief.pdf?sfvrsn=2ae4bcce_3&utm)

<sup>58</sup> David Watkins et al., "NCD Countdown 2030: Efficient Pathways and Strategic Investments to Accelerate Progress towards the Sustainable Development Goal Target 3.4 in Low-Income and Middle-Income Countries," *The Lancet* 399 (2022): 1266–78, <https://www.thelancet.com/journals/lancet/article/PIIS0140-67362102347-3/fulltext> Quoted in Ibid.

<sup>59</sup> Chisholm et al., "Scaling-up Treatment of Depression and Anxiety: A Global Return on Investment Analysis," *The Lancet Psychiatry* 3, no. 5 (2016): 415–24, [https://www.thelancet.com/journals/lanpsy/article/PIIS2215-0366\(16\)30024-4/fulltext](https://www.thelancet.com/journals/lanpsy/article/PIIS2215-0366(16)30024-4/fulltext), Quoted in Ibid.

<sup>60</sup> World Health Organization and World Bank Group (2024) 'International Dialogue on Sustainable Financing for Noncommunicable Diseases and Mental Health: report'. Available at [https://cdn.who.int/media/docs/default-source/ncds/sustainable-financing-dialogue-meeting-report.pdf?sfvrsn=40f650ff\\_4](https://cdn.who.int/media/docs/default-source/ncds/sustainable-financing-dialogue-meeting-report.pdf?sfvrsn=40f650ff_4)

<sup>61</sup> WHO and World Bank, June 2024, 'International Dialogue on Sustainable Financing for Noncommunicable Disease and Mental Health: Meeting report'.

<sup>62</sup> HelpAge International, 2017, How data systems leave older people behind.

<sup>63</sup> Saifur Rahman Chowdhury et al, 2023

<sup>64</sup> HelpAge, 2020, If not now, when? Keeping promises to older people affected by humanitarian crises

<sup>65</sup> WHO, 2022, Decade of Healthy Ageing Connection Series No. 3 - The Decade in a Climate-changing World

<sup>66</sup> NCD Alliance, 2024, [NCDs\\_neglected\\_crisis-humanitarian\\_brief-2024.pdf](#) (ncdalliance.org)

<sup>67</sup> NCD Alliance, 2024, NCD Alliance Advocacy Priorities for the 4th High-level Meeting of the UN General Assembly on the prevention and control of NCDs in 2025

<sup>68</sup> Ibid.

<sup>69</sup> Ibid.

<sup>70</sup> World Bank, 2024, Unlocking the Power of Healthy Longevity: Demographic Change, Non-communicable Diseases, and Human Capital (worldbank.org)

HelpAge International is a global network of organisations promoting the right of all older people to lead dignified, healthy and secure lives.

## Driving equitable action on NCDs and healthy ageing to achieve health for all at all ages

Published by HelpAge International  
PO Box 78840  
London  
SE1P 6QR  
UK

Tel +44 (0)20 7278 7778

info@helpage.org  
**www.helpage.org**

Registered charity no. 288180



Copyright © HelpAge International 2025

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License,  
<https://creativecommons.org/licenses/by-nc/4.0>

Any parts of this publication may be reproduced without permission for non-profit and educational purposes. Please clearly credit HelpAge International and send us a copy or link.